



DIRECTIONS FOR USE:

For the Control of Fly larvae:

Apply HokoEX at intervals exceeding 14 days. Animals may be present during applications. Often, 6-week intervals suffice.

Apply HokoEX at a dose of 250g/10m², independent of the method of application.

Apply HokoEX either as dry granules by dispersal if the manure is liquid or very moist, or dilute 250 g in 1-4 L of water for spraying or 10 L water for pouring using a watering can. In case the manure is dry apply by pouring only.

The simultaneous use of an insecticide specific to adult flies, particularly coinciding with the first application of HokoEX, can significantly enhance fly control.

In rearing facilities for CATTLE and HORSES apply HokoEX in a 50 cm band along the walls, under and next to machinery installations by pouring or spraying. When treating liquid manure the granules can be dispersed directly. The first application is best carried out 1-3 days after dung removal. In cow sheds with slatted floors apply HokoEX uniformly to the entire surface.

In PIG stys with completely slatted floors apply HokoEX to the entire surface, in the case of partially slatted floors apply to the slatted sections and the borders of the rest areas. Apply either by dispersal of dry granules, or pouring, or spraying. If possible treat before housing the pigs, shortly after manure or slurry removal is best.

In POULTRY houses with a slurry reservoir and in the presence of moist or liquid manure, disperse HokoEX granules directly, in dry manure situations use the pouring or spraying method. Apply to the entire surface. Commence treatment approximately 1 week after removal of excrement but before a depth of 10cm has accumulated.

Hokoex can also be used in sheep and goat sheds, rabbit farms, zoological gardens in the same manner.

25 g / m²



HokoEx[®]™

INSECTICIDE FOR PROFESSIONAL USE

HokoEX is a larvicide used to control fly larvae in animal husbandry. Contains a potent growth disruptor specific to all fly species.

CONTAINS:

Cyromazine:
Soluble Granule (SG) formulation.


2.0 % w/w (20 g/Kg)

MANUFACTURER:
Hokochemie GmbH
Niesenweg 4
CH-3012 Berne

Tel: + 41 31 302 84 04

Fax: + 41 31 302 84 10

www.hoko.com



Hokoex, Hoko and the figurative OKO mark are trade marks of Hokochemicals Ltd., Berne, Switzerland.

MARKETING COMPANY:

Hockley International Limited
Hockley House, 3 Longstone Road,
Ashbrook Office Park, Manchester M22 5LB,
United Kingdom.

Tel: + 44 (0) 161 209 7400

Fax: + 44 (0) 161 209 7401

www.hockley.co.uk

EXCLUSIVE DISTRIBUTOR:

Bowden & Knights,
Unit 6,
Dolphin Business Park,
Shadwell, Thetford,
Norfolk IP24 2RY,
United Kingdom

Tel: + 44 (0) 1953 681 830

Fax: + 44 (0) 1953 681 179

www.bowdenandknights.com



10 kg

USE PRECAUTIONS:

ASH ANY CONTAMINATION from skin and eyes immediately.
WASH HANDS AND EXPOSED SKIN before meals and after use.
Keep only in the original container. Keep container tightly closed. KEEP IN A SAFE PLACE. Keep out of the reach of children. Keep away from food, drink and animal feeding stuffs. COLLECT EGGS AND REMOVE FEED, EXPOSED WATER AND MILK BEFORE APPLICATION.
USE ONLY AS DESCRIBED ON THE LABEL.

STATUTORY CONDITIONS RELATING TO USE

FOR USE ONLY AS AN INSECTICIDE
FOR USE ONLY BY PROFESSIONAL OPERATORS

Dose: See Section "DIRECTIONS FOR USE" for details on dose rates

READ ALL PRECAUTIONS BEFORE USE. HSE NO. 8466

PRECAUTIONS

The (COSHH) Control of Substances Hazardous to Health Regulations 2002 may apply to the use of this product at work.
Engineering control of operator exposure must be used where reasonably practicable in addition to the following items of personal protective equipment.
Wear suitable protective clothing and gloves.
AVOID EXCESSIVE CONTAMINATION OF COVERALLS AND LAUNDRY REGULARLY. However engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.
DO NOT CONTAMINATE FOODSTUFFS, EATING UTENSILS OR FOOD CONTACT SURFACES.

This material and its container must be disposed of in a safe way.
DO NOT BREATHE DUST or spray mist. Otherwise wear respiratory protective equipment and eye protection (see HSE Guidance Booklet HS (G) 53: "Respiratory Protective Equipment - a practical guide for users".
COVER ALL WATER STORAGE TANKS before application.
DO NOT APPLY DIRECTLY TO LIVESTOCK AND POULTRY.
EXCLUDE ALL PERSONS DURING TREATMENT.

This product is approved under the Control of Pesticides Regulations 1986 (as amended) for use as directed.

READ ALL PRECAUTIONS BEFORE USE

HSE No. 8466

PCS No. 95098



USE ONLY IN ACCORDANCE WITH THE
DIRECTIONS ON THE LABEL